



Original Article



Epidemiological Analysis of Gastrointestinal Parasites in Various Breeds of Cattle in the Northern Region of Khyber Pakhtunkhwa, Pakistan

Maaz Saleem¹, Muhammad Zahid Shah¹, Abdul Jalil Khan¹, Sheeba Begum¹, Muhammad Hamza¹, Muhammad Farooq Khan¹, Farhan Ullah¹, and Iftikhar Ahmad^{1*}

¹Department of Basic and Applied Zoology, Shaheed Benazir Bhutto University, Sheringal Dir Upper, Pakistan

ARTICLE INFO

Keywords:

Cattle, Breeds, Gastrointestinal Parasite, Flotation, Sedimentation Technique

How to cite:

Saleem, M., Shah, M. Z., Khan, A. J., Begum, S., Hamza, M., Khan, M. F., Ullah, F., & Ahmad, I. (2025). Epidemiological Analysis of Gastrointestinal Parasites in Various Breeds of Cattle in the Northern Region of Khyber Pakhtunkhwa, Pakistan: Gastrointestinal Parasites in Various Breeds of Cattle in the Northern Region. MARKHOR (The Journal of Zoology), 6(1), 37-41. <https://doi.org/10.54393/mjz.v6i1.153>

*Corresponding Author:

Iftikhar Ahmad

Department of Basic and Applied Zoology, Shaheed Benazir Bhutto University, Sheringal Dir Upper, Pakistan
iftikhar@sbbu.edu.pk

Received Date: 8th February, 2025

Revised Date: 20th March, 2025

Acceptance Date: 27th March, 2025

Published Date: 31st March, 2025

ABSTRACT

Gastrointestinal (GI) parasites are major problem in cattle production worldwide. **Objectives:** To determine the common gastrointestinal parasites of cattle in district Lower Dir Khyber Pakhtunkhwa, Pakistan. **Methods:** A cross-sectional descriptive design was used. The cow fecal samples were taken and were examined under microscope by using the sedimentation and flotation techniques. To determine the impacts of breed, age, parity, eating habits, deworming status, and herd size. The data were subjected for statistical analysis via Chi-square test by using SPSS. **Results:** A total of (202/300), 67.3% of fecal samples were found positive for GI parasites. The parasites were more prevalent (41.6%, each) in the Friesian and Jersey breeds. Endo-parasitic infection was higher (92.6%; $p=0.04$) in female cattle than in male. Less than 2 years' cattle had a higher (75.2%; $p=0.101$) incidence of GI parasite. Cattle reared in a mixed feeding system had a higher (91.1%; $p=0.245$) incidence rate of parasitic infection. Herd of ≤ 5 animals had 72.3% infestation rate, and 6-8 animals per herd had 27.7% infestation. Cattle which are not treated with anthelmintic drugs had a higher (53%; $p=0.988$) incidence rate of GI parasites than those that are treated with anthelmintic drugs (47%). *Taxocara vitulorum* prevalence was high (46%) in single parasitic infestation, while *Taxocara vitulorum* + *Haemonchus contortus* were higher (34.6%) in double parasitic infection. *Haemonchus* + *Taxocara* + *Fasciola* spp were detected more (33.3%) in triple parasitic infection. **Conclusions:** It was concluded that the cattle population in district lower Dir had the highest prevalence of gastrointestinal parasites and need effective control measures to enhance productivity.

INTRODUCTION

Livestock play a crucial role in the economy of Pakistan, with a share of 62.68% in agriculture value added and 14.36% in GDP [1]. Almost eight million families in Pakistan with 30-35 million rural populations are involved in livestock production activities and derive 35-40% of their annual income from this sector [2]. Livestock is a crucial asset, a key source of food and a potential source of income for the poor and landless farmers in Khyber Pakhtunkhwa and has a significant role in the provincial economy through its contribution of 57.5 % to the GNP. However, diseases and parasites are among the major constraints that impact livestock productivity, particularly for small-scale farmers,

due to poor disease diagnosis and limited understanding of the risk factors, hence affecting food supplies, commerce trade, and human health. Among parasites, the gastrointestinal parasites of different genera that inhabit the digestive tract of cattle, sheep and goats cause inappetence, poor feed utilization, anaemia, diarrhea, poor growth, decreased milk production, morbidity, mortality hence huge economic losses to the industry [3, 4]. In Pakistan, the prevalence of the parasitic infestation is very common and causes enormous losses to livestock [5, 6]. It has been observed that the prevalence of gastrointestinal helminthes is associated with agroclimatic conditions



such as temperature, humidity, rainfall, quantity and quality of the feed and grazing behavior of animals [7]. Gastrointestinal parasites have been extensively studied in Pakistan with variable results concerning the type of parasite, species of animal, agroclimatic conditions and season [8, 9]. Hence, it is of prime importance to identify the burden and types of helminthes and the associated risk factors for specific areas for the effective control measures. Information on the prevalence of gastrointestinal parasites is scarce in different livestock species, available in the districts of Lower Dir and Upper Dir [10, 11]. However, a precise investigation on the prevalence of gastrointestinal parasites and the associated risk factors is lacking in the study area.

This study aimed to explore the prevalence of gastrointestinal parasites and the associated risk factors to provide widespread information for designing effective control measures.

METHODS

A cross-sectional descriptive designed study was conducted at the district Lower Dir (KP) [13]. The district is located between 1200 and 2800 meters above sea level in the northern KP highlands, with latitudes of 34.35 and longitudes of 71.85. The Open Epi program was used to determine the sample size. The average annual rainfall was 1186 mm, and the average annual temperature was 16°C. During the wet and dry seasons, the relative humidity ranges from 70 to 81% and 40 to 50%, respectively. One fecal sample (~50 g) per cattle was directly collected from the rectum with gloved hands. The samples were immediately transferred to pre-labelled, hygienic plastic bottles. The sampling bottles were stored in screw capped container with ice packs and transported to laboratory. All samples were analyzed for the detection of parasites within 24 h of collection. Data on breed, sex, age, herd size, deworming status and feeding pattern, date and place of sampling were recorded for each sampled animal at the time of sampling. fecal samples were processed and examined by direct and indirect parasitological techniques (centrifugation, flotation and sedimentation) [12]. The GI parasites were identified using identification keys described by Otranto D and Wall R [13]. Briefly, fecal materials (1 g) were mixed with 0.9% normal saline wet mount solution in a mortar and a relatively homogenous preparation was obtained. The suspension was then filtered through a tea strainer. Finally, a drop of suspension was added to a glass slide and examined under a microscope. From each fecal sample, three direct smears were examined. For flotation technique, two grams of feces was put in sterile screw capped bottle, containing 5 ml of the saline solution and was mixed and strained through a sieve. The mixed matters were riddled into a centrifuge tube or a walled test tube. Formalin was added to the test

tube until a convex meniscus was formed. A cover slip was positioned on the top of the test tube carefully and left for 5 minutes. The cover slip was removed from the glass tube and placed on the slide, and was tested for helminthes eggs and oocysts under the microscope at 10X. In the sedimentation method, two g of feces samples were mixed with 50 ml of water and sieved. The suspension was centrifuged at 1000 rpm for 2-3 minutes with NaCl solution. The supernatant was decanted, and from the sediment, 1 drop was taken with a Pasteur's pipette and put on a slide and examined under the microscope at 10X for the presence of helminthes eggs. The data were presented in percentiles. Statistical Package for Social Sciences (SPSS) version 23.0 was used for data analysis, using the Chi-square test.

RESULTS

According to the present study, a total 67.3% samples were positive for parasitic infestation. Breed-wise analysis indicates highest (41.6% each breed) prevalence in Friesian and Jersey cow. Values are presented in percentages. The percentage value has been calculated from the total number of fecal samples examined (n=300). Data showing the overall prevalence of parasitic infestation in various breeds of cattle are presented in table 1.

Table 1: Prevalence (%) of Gastrointestinal Parasites in Various Breeds of Cattle (n=300)

Breed	Sample Size	Positive	Prevalence	p-Value
Friesian	134	84	41.6%	0.116
Jersey	111	84	41.6%	
Achai	39	25	12.4%	
Non-descript	16	9	4.5%	
Total	300	202	67.3%	

Female cattle had a high (92.6%) prevalence of GI parasites. Cattle less than 2 years old had a high (75.2%) prevalence of GI parasites. Likewise, cattle reared under a mixed-type feeding system had a higher (91.1%) incidence of GI parasites than cattle under stall feeding. Notably, 72.3% of samples were detected positive in cattle with a herd size of ≤5 animals, while 6-8 animals per herd had 27.7% positive samples for GI parasites. Cattle having no history of anthelmintic use had a 53% detection rate of parasitic infection. Those partially dewormed had a 47% detection rate (Table 2).

Table 2: The Effect of Sex, Age, Feeding, Herd Size and Deworming Status on the Prevalence of GI Parasites in Cattle Breed

Breed	Sample Size	Positive	Prevalence	p-Value
Gender				
Male	17	15	7.4%	0.04
Female	283	187	92.6%	

Age (Years)				
<2	214	152	75.2%	0.101
3-4	76	45	22.3%	
5-6	9	4	2%	
>6	1	1	0.5%	
Feeding				
Mixed Feeding	277	184	91.1%	0.245
Stall Feeding	23	18	8.9%	
Herd Size				
≤ 5	217	146	72.3%	0.975
6-8	83	56	27.7%	
Deworming				
Partially Deworm	141	95	47%	0.988
No	159	107	53%	

Data is in percentage and has been calculated from the total number of positive faecal samples observed (n=202)

Among the reported gastrointestinal nematodes, *Toxocara vitulorum* had a high (46%) prevalence rate, followed by *Haemonchus contortus* (19%), then other parasites. Among the gastrointestinal cestodes, *Moniezia spp* had a 17% prevalence rate. In the trematode parasite, *Fasciola hepatica* had 12% positive cases. Importantly, *Toxocara vitulorum* + *haemonchus contortus* were more (34.6%) prevalent in double parasitic infection. While *Haemonchus* + *Taxocara* + *Fasciola spp* were detected more (33.3%) in triple parasitic infection. The species-wise prevalence of parasites in various breeds of cattle is shown in table 3.

Table 3: Prevalence of Single, Double and Triple Parasites Species in Cattle (n=200)

Parasite Species		n (%)
Nematodes	<i>Toxocara vitulorum</i>	46 (46%)
	<i>Haemonchus contortus</i>	19 (19%)
	<i>Trichuris</i>	2 (2%)
	<i>Monezia spp</i>	17 (17%)
	<i>Dictyocaulus viviparus</i>	1 (1%)
	<i>Ostertaigia</i>	1 (1%)
Trematode	<i>Fasciola hepatica</i>	12 (12%)
	<i>Trichostrongylus</i>	1 (1%)
Protozoan	<i>Eimeria bovis</i>	1 (1%)
Double Infection of Parasitic Species	<i>Taxocara vitulorum</i> + <i>Haemonchus contortus</i>	27 (34.6%)
	<i>Taxocara vitulorum</i> + <i>Fasciola hepatica</i>	12 (15.4%)
	<i>Haemonchus contortus</i> + <i>Fasciola hepatica</i>	11 (14.1%)
	<i>Taxocara vitulorum</i> + <i>Monezia spp</i>	8 (10.3%)
	<i>Monezia spp</i> + <i>Fasciola hepatica</i>	8 (10.3%)
	<i>Haemonchus contortus</i> + <i>Monezia spp</i>	7 (9.0%)
	<i>Taxocara vitulorum</i> + <i>Trichuris</i>	3 (3.8%)
	<i>Haemonchus contortus</i> + <i>Eimeria bovis</i>	2 (2.6%)
Mixed Infection of Parasitic Species	<i>Taxocara vitulorum</i> + <i>Haemonchus contortus</i> + <i>Fasciola hepatica</i>	8 (33.3%)
	<i>Taxocara vitulorum</i> + <i>Monezia spp</i> + <i>Fasciola hepatica</i>	5 (20.8%)
	<i>Taxocara vitulorum</i> + <i>Haemonchus contortus</i> + <i>Monezia spp</i>	3 (12.5%)

<i>Taxocara vitulorum</i> + <i>Haemonchus contortus</i> + <i>Teania</i>	2 (8.3%)
<i>Haemonchus contortus</i> + <i>Monezia spp</i> + <i>Fasciola hepatica</i>	2 (8.3%)
<i>Taxocara vitulorum</i> + <i>Trichuris</i> + <i>Fasciola hepatica</i>	1 (4.2%)
<i>Haemonchus contortus</i> + <i>Fasciola hepatica</i> + <i>Ostertegia</i>	1 (4.2%)
<i>Trichuris</i> + <i>Haemonchus contortus</i> + <i>Moniezia spp</i>	1 (4.2%)
<i>Haemonchus contortus</i> + <i>Moniezia spp</i> + <i>Eimeria bovis</i>	1 (4.2%)

The microscopic appearance of different parasitic species is shown in figure 1.




Figure 1: Different Parasitic Species

DISCUSSION

Parasitism is one of the most serious issues that the cattle population faces around the world. Parasitic diseases, particularly gastrointestinal nematode and trematode infections, pose a severe health risk to cattle and reduce output due to related morbidity, mortality, treatment costs, and control measures [14]. A thorough grasp of the disease epidemiology, pasture management, farm management techniques, and agroclimatic factors like rainfall and temperature is all necessary for the control of gastrointestinal parasite infections in animals [15]. The kind of parasite, the extent of the infestation, and additional risk variables such as species, age, and season

all are responsible for gastrointestinal parasite infections [16]. According to the present study, a total of 67.3% samples were positive for parasitic infestation. Breed-wise analysis indicates the highest (41.6% each breed) prevalence in Friesian and Jersey cows. In the previous study, Endo-parasites were detected in 43.96% of the total fecal samples, which was less than the 47.00% of Nigerian cattle housed at the study farm [17]. According to our findings, nematodes (66.99%) were the most common helminthes infection, followed by cestodes (17%), and tally trematode (13%). The parasite infection ratio is only found in one protozoan (3%). In a prior study, nematodes (72.41%) were the most common helminth infection, followed by trematodes (25.00%) and cestodes (25.00%). Strongyles were found in larger numbers in cattle [17]. According to our analysis, parasite infestation was highest in the Friesian cross (76.6%), followed by the Jersey cross (68.6%), Sahiwal (50%) and Achai (48.7%). In the nondescript, the parasite infection was the lowest. A significant difference ($p < 0.05$) was also recorded with the infestation levels in different breeds in the prior study, local breed cattle having a higher infection rate than cross-breed cattle. In comparison to reports of GI nematode infection in cross-breed cattle, the prevalence of GI nematode infection in cross-breed cattle was lower. According to our analysis, parasitic infection was most common in animals aged 6 to 10, followed by animals aged 1 to 5, and finally, animals aged 16 to 20. The animals in the age group 11-15 had the lowest prevalence. The frequency of GI nematode infection was found to be higher in comparison to a previous study in those animals aged less than 1 year on Haramaya University dairy farm on the Holstein Friesian dairy breed followed by juvenile and mature [18]. According to our analysis, females have the highest parasite infestation followed by males. Furthermore, they had a larger endo-parasite infection percentage of *Haemonchus contortus* than males, according to [19]. The higher prevalence of gastrointestinal parasites could be linked to cattle management. The mixed feeding group had the highest parasitic infestation in the current study's grazing pattern. The animals with the 1-4 herd size had the highest prevalence ratio in the herd size category. Among terms of deworming techniques, the highest parasitic infestation was found in animals that had never been dewormed in their whole lives [20]. The lowest frequency rate of gastrointestinal parasites is due to deworming and care techniques. The majority of cattle are untreated, while others were grazing animals, who were rarely treated for GI diseases. Grazing animals have more chances for entry of various parasite stages into cattle's digestive tracts via oral ingestion.

CONCLUSIONS

It was concluded that numerous internal parasites are prevalent in cattle, with a greater infection incidence in the Friesian and Jersey crossbreed. Infestation of parasitic nematodes was high. On the other hand, the highest prevalence was recorded in female. In the age the highest prevalence was recorded in the age group of 6-10 years. Mixed feeding pattern, parasite infestation was common in diverse management approaches. The highest prevalence was recorded in those cattle which have parity level. The highest prevalence was recorded in those animals which do not properly dewormed. High parasite infestation were recorded in the small herd size as most of the people had a small herd size in the study area from where samples were collected. Risk factors had a close relation with parasite infestation.

Authors Contribution

Conceptualization: MS, IA

Methodology: MS, IA

Formal analysis: MH, MFK, FU

Writing review and editing: MZS, AJK, SB

All authors have read and agreed to the published version of the manuscript.

Conflicts of Interest

All the authors declare no conflict of interest.

Source of Funding

The author received no financial support for the research, authorship and/or publication of this article.

REFERENCES

- [1] Khan T, Khan W, Iqbal R, Maqbool A, Fadladdin YA, Sabtain T. Prevalence of Gastrointestinal Parasitic Infection in Cows and Buffaloes in Lower Dir, Khyber Pakhtunkhwa, Pakistan. *Brazilian Journal of Biology*. 2022 Feb; 83: e242677. doi: 10.1590/1519-6984.242677.
- [2] Abbas N, Qayyum M, Hasan M, Shoaib M, Zafar A, Riaz A. Prevalence and Epidemiology of Gastrointestinal Parasites in Cattle in Different Zones of Tehsil Chakwal, Punjab. Pakistan. *Journal of Bacteriology and Parasitology*. 2021; 12(8).
- [3] León JC, Delgado NU, Florez AA. Prevalence of Gastrointestinal Parasites in Cattle and Sheep in Three Municipalities in the Colombian North-eastern Mountain. *Veterinary World*. 2019 Jan; 12(1): 48. doi: 10.14202/vetworld.2019.48-54.
- [4] Tachack EB, Oviedo-Socarrás T, Pastrana MO, Pérez-Cogollo LC, Benavides YH, Pinto CR et al. Status of Gastrointestinal Nematode Infections and Associated Epidemiological Factors in Sheep from

- Córdoba, Colombia. *Tropical Animal Health and Production*. 2022 Jun; 54(3): 171. doi: 10.1007/s11250-022-03170-2.
- [5] Rizwan HM, Zohaib HM, Sajid MS, Tahir UB, Kausar R, Nazish N et al. Unveiling the Hidden Threat: Investigating Gastrointestinal Parasites and Their Costly Impact on Slaughtered Livestock. *Revista Brasileira de Parasitologia Veterinária*. 2024 Oct; 33(3): e007224. doi: 10.1590/s1984-29612024061.
- [6] Rizwan HM, Zohaib HM, Sajid MS, Abbas H, Younus M, Farid MU et al. Inflicting Significant Losses in Slaughtered Animals: Exposing the Hidden Effects of Parasitic Infections. *Pathogens*. 2023 Oct; 12(11): 1291. doi: 10.3390/pathogens12111291.
- [7] Terfa W, Kumsa B, Ayana D, Maurizio A, Tessarin C, Cassini R. Epidemiology of Gastrointestinal Parasites of Cattle in Three Districts in Central Ethiopia. *Animals*. 2023 Jan; 13(2): 285. doi: 10.3390/ani13020285.
- [8] Shah SS, Khan MI, Ullah A, Ullah H, Ahmad F. Coprological Examination of Small and Large Ruminants in Central Zone of Khyber Pakhtunkhwa. *Sarhad Journal of Agriculture*. 2021 Feb; 37(1): 152-7. doi: 10.17582/journal.sja/2021/37.1.152.157.
- [9] Lateef M, Akhtar R, Zahid B, Durrani UF, Qadry A, Afzal N et al. Prevalence of *Trichostrongylus* in Sheep in the District Zhob, Balochistan, Pakistan. *Arquivo Brasileiro de Medicina Veterinária e Zootecnia*. 2021 May; 73(02): 522-4. doi: 10.1590/1678-4162-12187.
- [10] Ruhollah, Khan W, Al-Jabr OA, Khan T, Khan A, El-Ghareeb WR et al. Prevalence of Gastrointestinal Parasite in Small Ruminants of District Dir Upper Khyber Pakhtunkhwa Province of Pakistan. *Brazilian Journal of Biology*. 2021 Oct; 83: e248978. doi: 10.1590/1519-6984.248978.
- [11] Khan T, Nasreen N, Shater AF, Khan A, Kamal M, Vinueza R et al. Risk Factor Analysis for the Prevalence of Gastrointestinal Parasites Found in Large Ruminants in Lower Dir Khyber Pakhtunkhwa Pakistan. *Saudi Journal of Biological Sciences*. 2021 Dec; 28(12): 7022-6. doi: 10.1016/j.sjbs.2021.07.078.
- [12] Soulsby EJ. *Helminths, Arthropods and Protozoa of Domesticated Animals*. 1982.
- [13] Otranto D and Wall R. *Veterinary Parasitology*. John Wiley and Sons. 2024 Mar. doi: 10.1002/9781394176373.
- [14] Shoaib M. Prevalence of the Gastrointestinal Helminths in Bovine Population in Different Zones of Rawalpindi District of Punjab, Pakistan. *Pakistan Journal of Science*. 2016; 68(4). doi: 10.57041/pjs.v68i4.245.
- [15] Gunathilaka N, Niroshana D, Amarasinghe D, Udayanga L. Prevalence of Gastrointestinal Parasitic Infections and Assessment of Deworming Program among Cattle and Buffaloes in Gampaha District, Sri Lanka. *Bio-Med Research International*. 2018; 2018(1): 3048373. doi: 10.1155/2018/3048373.
- [16] Tiele D, Sebro E, H/Meskel D, Mathewos M. Epidemiology of Gastrointestinal Parasites of Cattle in and Around Hosanna Town, Southern Ethiopia. *Veterinary Medicine: Research and Reports*. 2023 Dec; 1-9. doi: 10.2147/VMRR.S389787.
- [17] Nyutu CW. Prevalence of Gastrointestinal Parasites of Cattle in Mathira Constituency, Kenya (Doctoral Dissertation, Karatina University). 2021.
- [18] Hiko A and Wondimu A. Occurrence of Nematodiasis in Holstein Friesian Dairy Breed. *The Journal of Veterinary Medicine and Animal Health*. 2011 Jan; 3(1): 6-10.
- [19] Khan A, Jamil M, Ullah S, Ramzan F, Khan H, Ullah N et al. The Prevalence of Gastrointestinal Nematodes in Livestock and Their Health Hazards: A Review. *World's Veterinary Journal*. 2023; 2023(1): 57-64. doi: 10.54203/scil.2023.wvj6.
- [20] Piratheepan S. Cross-Sectional Study of Gastrointestinal Parasitism of Cattle in Kilinochchi District of Sri Lanka. *Veterinary Practitioner*. 2024 Dec; 25(2).